

TWISTOR ALGEBRAIC DYNAMICS IN COMPLEX SPACE-TIME AND PHYSICAL MEANING OF HIDDEN DIMENSIONS

Vladimir V. Kassandrov

Abstract. We review the algebraic field theory based completely on a nonlinear generalization of the CR complex analyticity conditions to the noncommutative algebra of biquaternions. Any biquaternionic field possesses natural twistor structure and, in the Minkowski space, gives rise to a shear-free null congruence of rays and to a set of gauge fields associated with it. In the paper we develop the algebrodynamical scheme in the complex extension of Minkowski space-time – in the full vector space of biquaternion algebra. This primordial space dynamically reduces to the 6D “observable” space-time of the complex null cone which in turn decomposes into the 4D physical space-time and the 2D internal “spin space”. A set of identical point charges (“duplicons”) – focal points of the congruence – arises in the procedure. Temporal dynamics of individual duplicons is strongly correlated via fundamental twistor field of the congruence. We briefly discuss some new notions inevitably arising in the considered algebrodynamical scheme, namely those of “complex time” and of “evolutionary curve”, as well as their hypothetical links to the quantum uncertainty phenomenon.

1 Introduction

In the article we continue to develop the *algebrodynamical* field theory which has been presented, in particular, in [1, 2]. We recall that in the paradigm of algebrodynamics one considers, in the spirit of radical Pithagorean philosophy, all fundamental physical laws and phenomena as a manifestation of some basic abstract “World” structure – *the Code of the Universe* – which might be an exceptional group (e.g., the *Monster group*), algebra or geometry. The whole structure and evolution of the Universe, the categories of time, particle, field, motion and interaction should be all predetermined and encoded in internal properties of the World structure. The known physical laws are nothing but separate *fragments* of this basic structure, and one could try to *forget* them all and to read *unprejudicely* the “Book of Nature” which could bring us by this to a grand and magestic picture of physical World *quite different from that accepted in the modern theoretical physics*.

In the main version of algebrodynamics developed by the author (see, e.g., [3, 4, 5, 6, 7] and references therein) the exceptional algebra of *quaternions* (precisely, of their complexification – the algebra of *biquaternions* $\mathbb{B}$) is considered as the *World algebra*. Basic physical fields are then $\mathbb{B}$ -valued functions of $\mathbb{B}$ -variable, and field equations are nothing but the *conditions of $\mathbb{B}$ -differentiability*, i.e. the *generalized Cauchy-Riemann equations* (GCRE). These have been introduced by the author in 1980 (see in detail [3, 4, 8, 9, 10]) and have the following invariant form:

$$dF = \Phi * dZ * \Psi, \quad (1)$$

where $F : \mathbb{B} \mapsto \mathbb{B}$ is a $\mathbb{B}$ -differentiable function $F(Z)$ of $\mathbb{B}$ -variable $Z \in \mathbb{B}$; $\Phi, \Psi : \mathbb{B} \mapsto \mathbb{B}$ are two auxiliary functions (left and right “semiderivatives”) associated with F , and $(*)$ denotes the operation of multiplication in $\mathbb{B}$ isomorphic to the full 2×2 complex matrix algebra.

As a direct consequence of (1), any matrix component of the $\mathbb{B}$ -field $f = \{F_{AB}\}$, $A, B = 1, 2$ obeys the determinant-type equation

$$\det \left\| \frac{\partial f}{\partial Z_{AB}} \right\| = 0 \quad (2)$$

which, for the case of noncommutative $\mathbb{B}$ -algebra, is a *nonlinear* analogue of the Laplace *harmonicity* equation in complex analysis.

Let us restrict now the coordinates Z onto the subspace $\mathbf{M}$ of the full $4\mathbb{C}$ vector space of $\mathbb{B}$ -algebra represented by Hermitian matrices $X = X^+$. This has the metrical structure identical to that of *Minkowski* space. Below we'll use the following representation for the coordinates on $\mathbf{M}$:

$$Z \mapsto X = \begin{pmatrix} u & w \\ p & v \end{pmatrix} \quad (3)$$

where $u = ct + z$, $v = ct - z$ are real, $w = x - iy$ and $p = x + iy$ – complex conjugated and x, y, z, t – Cartezian and time coordinates respectively. Now the GCRE read as follows:

$$dF = \Phi * dX * \Psi, \quad (4)$$

and the fundamental equation (2) takes the form

$$\frac{\partial f}{\partial u} \frac{\partial f}{\partial v} - \frac{\partial f}{\partial w} \frac{\partial f}{\partial p} \equiv \frac{1}{c^2} \left(\frac{\partial f}{\partial t} \right)^2 - \left(\frac{\partial f}{\partial x} \right)^2 - \left(\frac{\partial f}{\partial y} \right)^2 - \left(\frac{\partial f}{\partial z} \right)^2 = 0 \quad (5)$$

in which one easily recognizes the *complex eikonal equation* (CEE) [3].

On the Minkowski coordinate background the whole theory turns to be Lorentz invariant, and one is able to construct an “algebraic” field theory in which the nonlinear complex eikonal is a basic *self-interacting* physical field.

Principles and results of (nonlinear, non-Lagrangian, overdetermined) algebraic $\mathbb{B}$ -field theory have been presented in a series of papers (see, e.g., [1, 2, 4, 9, 10, 11, 12, 13, 7] and other papers). The GCRE and the related CEE were found to possess 2-spinor, twistor and restricted gauge structures. Moreover, *the integrability conditions for (4) result in identical satisfaction of the (complexified) homogeneous Maxwell and Yang-Mills equations.*

In our paper [13] the Lorentz invariant form of *general solution* to the CEE has been found on the base of its (ambi)twistor structure. Precisely, any (almost everywhere analytical) solution of CEE can be obtained from some generating function

$$\Pi(G, \tau^1, \tau^2) \equiv \Pi(G, wG + u, vG + p) \quad (6)$$

of three complex variables – components of the projective (null) twistor related to the points of space-time $X \in \mathbf{M}$ via the Penrose *incidence condition* [14]

$$\tau = X\xi \quad (\tau^A = X^{AA'}\xi_{A'}), \quad (7)$$

from which, under the choice of the gauge $\xi_{A'}^T = \{1, G\}$, one gets the constraints

$$\tau^1 = wG + u, \quad \tau^2 = vG + p. \quad (8)$$

Any generating twistor function (6) gives rise to a pair of the CEE solutions. To obtain the first one, one resolves the algebraic constraint

$$\Pi(G, wG + u, vG + p) = 0 \quad (9)$$

with respect to G and comes then to a complex (and a *multivalued* in general) field $G(u, v, w, p)$ (or $G(x, y, z, t)$) which *for any* Π *identically satisfies the CEE*. Secondly, one differentiates the function (6), resolves then again the arising constraint

$$P \equiv \frac{d\Pi}{dG} = 0 \quad (10)$$

with respect to G and substitutes the resulting field $G(u, v, w, p)$ into (6). In this way one comes to a function of coordinates $\Pi(G(X), \tau^1(X), \tau^2(X)) \equiv \Pi(x, y, z, t)$ which again *identically satisfies the CEE*. It was shown in [13] that the two classes above presented *exhaust the solutions to CEE*.

Geometrically, any twistor field $\mathbf{W}(X) = \{G, \tau^1, \tau^2\}$ defines a *null geodesic congruence*, i.e. a congruence of rectilinear light-like rays on $\mathbf{M}$. For twistors obtained from the constraint (9) this congruence is known to be always *shear-free* (SFC) and, moreover, any SFC can be constructed in this way (this is the content of the so called *Kerr theorem* [14, 15]).

Conditions (9),(10) considered together specify the *caustic locus* of the related congruence so that after elimination of G the resulting equation

$$\Pi(u, v, w, p) \equiv \Pi(x, y, z, t) = 0 \quad (11)$$

(with the function Π always obeying the CEE, see above) *determines the shape and the temporal evolution of caustics* which can be enormously complicated [16, 6, 2]. On the other hand, the same condition defines the *singular locus* of the associated Maxwell and Yang-Mills fields [11, 9], as well as of the curvature field of correspondent Kerr-Schild metrics [15, 17]. Therefore, we can identify *particles as singularities*¹, *common to the electromagnetic and the other fields defined by any solution of the GCRE (or, equivalently, by any SFC)* [16, 11, 9].

Note that these singularities possess some of the properties of real quantum particles due, in particular, to the *overdetermined* structure of the GCRE. In fact, according to the *theorem of charge quantization* proved in [12, 6], any bounded and isolated singularity of electromagnetic field associated with solutions of GCRE has the value of electric charge either zero or integer multiple to some minimal, *elementary* one, precisely to the charge of static axisymmetrical *Kerr solution* with a ring-like singularity [3, 4, 11, 10]. According to observations of Carter [18] and Newman [19], the gyromagnetic ratio for the Kerr-like singularities is exactly the same as that for Dirac fermions. Numerous examples of SFC with complicated structure and dynamics of singular loci has been presented in our works [11, 16, 10, 9, 2].

From general viewpoint, we come in this way to a peculiar picture of physical World whose main elements are the congruence of the “primordial light” rays (invisible for an ordinary observer) – the flow of *Prelight* [24, 1, 2] – and the matter “born” from this “Ether-like” yet Lorentz invariant flow at its focal points, i.e. at caustics. A consistent concept of physical Time as the *field evolutional parameter* along the rays of the Prelight flow has been also developed and discussed in our papers [1, 2, 24, 25] (see also section 3).

Unfortunately, the above presented algebraic theory for interacting fields-particles is far from being realistic. Indeed, generically therein the particles-caustics are represented

¹At least for the case when the singular locus is bounded in 3D space

by a number of isolated 1D curves (*strings*) nontrivially evolving in time but, as a rule, unstable in shape and size and radiating to infinity. Besides, one finds no correlation in temporal dynamics of different strings which could model their interaction. Finally, from the epistemological viewpoint, the real Minkowski subspace is by no means distinguished in the structure of $\mathbb{B}$ -algebra: its elements doesn't even form a subalgebra of $\mathbb{B}$.

The last argument is of an especial importance in the framework of the algebrodynamics. In fact, no algebra is known whose *automorphism group* is isomorphic to the Lorentz group of Special Relativity. Therefore, in the algebrodynamical paradigm one is forced to admit space-time geometries different from the Minkowski one. By this, the full 4D *complex* vector space of the $\mathbb{B}$ -algebra seems to be the most appropriate background for the algebraic field-particle theory based on the $\mathbb{B}$ -differentiability conditions, i.e. on the GCRE.

Complex extension $\mathbf{CM}$ of the Minkowski space-time indeed arises repeatedly in General Relativity, in twistor and string theories. In particular, Newman et al. [19, 20, 21] made use of the concept of complex space-time in order to obtain physically interesting solutions of electrovacuum Einstein-Maxwell system and to link together physical characteristics of corresponding particle-like singular sources (see the next section). In our recent paper [25] the algebrodynamics in the $\mathbf{CM}$ space has been developed on the base of the Newman's representation for "virtual" point charge "moving" along a complex world line in $\mathbf{CM}$ and generating therein a congruence of complex "light-like rays". The "cut" of this null congruence by the real Minkowski slice $\mathbf{M}$ gives there rise to a SFC with interesting physical properties, in particular with nonzero *twist*.

Here again, however, one deals with a rather artificial and insufficient restriction of the basic structures onto $\mathbf{M}$. Alternatively, in the framework of the algebrodynamics in $\mathbf{CM}$ [25], we have introduced the concept of the *observable* space-time – the 6D subspace of the *complex null cone* (CNC) of a point-like "observer" $\mathbf{O}$ in $\mathbf{CM}$. By this, any particle-like (caustic) element $\mathbf{C}$ which could be detected by $\mathbf{O}$ *necessarily lies on its CNC* and has the same value of the primary twistor field as in $\mathbf{O}$, so that the temporal dynamics of $\mathbf{C}$ and of $\mathbf{O}$ are strongly correlated (they "interact").

On the other hand, the CNC has the topology $\mathbb{R}_+ \times \mathbf{S}^3 \times \mathbf{S}^2$ (see section 3), so that it decomposes into a basic 4D space (in which the time interval should be identified with the 4D distance as in [26, 27]) and an orthogonal space of a 2-sphere which can be naturally treated as an internal *spin space*.

Moreover, we'll see that an ensemble of identical point-like particles – *duplicons* – images of one and the same generating charge with correlated temporal evolution – can be naturally constructed in this framework. We'll also discuss probable physical sense of the *quaternionic time*, *complex time* and of the *evolutionary curve* – of notions inevitably arising in the considered algebrodynamical scheme and related, perhaps, to the origin of quantum uncertainty.

2 Newman's charge in $\mathbb{CM}$ and the set of its images – duplicons

Simplest variety of a shear-free null congruence (SFC) is formed by a bundle of light-like rays from a point particle moving along an arbitrary world line in the real $\mathbf{M}$ space [31]. Electromagnetic field $F_{\mu\nu}$ can be then associated with such a SFC via integration of Maxwell equations with additional requirement on the field strength $F_{\mu\nu}k^\nu = 0$ to be orthogonal to the null 4-vector $k^\mu = \xi_A \xi_{A'}$ tangent to the congruence rays. Quite expectably, this field is just the Lienard-Wichert one. In the algebrodynamical approach, the field of exactly the same type can be obtained directly via second derivatives of the congruence field G [4, 11, 10]); however, this field necessarily corresponds to an *elementary (unit) value of generating charge*². Notice that in the static case the SFC generated by the charge at rest is radial and gives rise to the “quantized” Coulomb field and to the Reissner-Nördstrom metrics.

Newman et al. proposed [23, 22] a generalization of this construction via consideration of a point-like charge “moving” in the complex extension $\mathbb{CM}$ of $\mathbf{M}$ (in fact, even the case of the *curved* space-time was studied). Then on the real slice $\mathbf{M}$ of $\mathbb{CM}$ one obtains a more complicated SFC with nonzero *twist* which, in the particular case of “virtual” charge at rest, gives rise to the metric and electromagnetic field of the well-known *Kerr-Newman* electrovacuum solution with a *ring-like singularity* (“Kerr’s ring”).

Consider now the Newman’s construction in twistor terms [32, 33] and its generalization in the framework of algebrodynamics. Let a point singularity “moves” along a complex world line $Z_\mu = \hat{Z}_\mu(\tau)$, $\mu = 0, 1, 2, 3$ where the parameter $\tau \in \mathbb{C}$ plays the role of “complex time” (further on we’ll consider its own “evolutionary law” $\tau = \tau(s)$, $s \in \mathbb{R}$). For Cartezian and spinor coordinates on $\mathbb{CM}$ we’ll use the following representation similar to (3):

$$Z = \{Z^A_B\} = \begin{pmatrix} u & w \\ p & v \end{pmatrix} = \begin{pmatrix} z_0 - iz_3 & -iz_1 - z_2 \\ -iz_1 + z_2 & z_0 + iz_3 \end{pmatrix} \quad (12)$$

in which all of the z_μ (as well as u, v, w, p) are now complex-valued. Incidence relation (7) takes then the form

$$\tau = Z\xi \quad (\tau^A = Z^A_B \xi^B, \quad \tau^1 = wG + u, \quad \tau^2 = vG + p), \quad (13)$$

where the quantities $\{G, \tau^1, \tau^2\}$, $G \equiv \xi^2/\xi^1$ constitute the projective twistor in $\mathbb{CM}$. If now a point $Z \in \mathbb{CM}$ is separated from the charge $\hat{Z}$ by the null interval,

$$S := \det \|Z - \hat{Z}(\sigma)\| = (u - \hat{u}(\sigma))(v - \hat{v}(\sigma)) - (w - \hat{w}(\sigma))(p - \hat{p}(\sigma)) = 0, \quad (14)$$

then one can find from (14) the field $\sigma = \sigma(Z) = \sigma(u, v, w, p)$ and, consequently, the position of the charge $\hat{Z}(Z) \equiv \hat{Z}(\sigma(Z))$ which “influences” the point Z . This means that *twistor field* is the same both at Z and at $\hat{Z}$. Indeed, in account of (14) the linear system of equations

$$(Z - \hat{Z}(\sigma))\xi = 0 \quad (15)$$

²Owing to the overdetermined structure of the GCRE (or of SFC equations), see section 1

in a unique way fixes the ratio of spinor components $G = \xi^2/\xi^1$. System (15) then reads as

$$\tau \equiv Z\xi = \widehat{Z}\xi, \quad (16)$$

so that all of the twistor components are indeed equal at Z and $\widehat{Z}$.

Notice that the complex parameter σ , as a function of the point of observation Z , always satisfy the eikonal equation. Now in components system (15) takes the form

$$\begin{cases} (u - \widehat{u}(\sigma)) + (w - \widehat{w}(\sigma))G &= 0 \\ (p - \widehat{p}(\sigma)) + (v - \widehat{v}(\sigma))G &= 0 \end{cases} \quad (17)$$

and demonstrates that the field G is *indefinite* on the world line of the charge itself (and depends on the direction in its vicinity). On the other hand, writing out (16) in the form

$$\begin{cases} \tau^1 &= \widehat{u}(\sigma) + \widehat{w}(\sigma)G \\ \tau^2 &= \widehat{p}(\sigma) + \widehat{v}(\sigma)G \end{cases} \quad (18)$$

and eliminating from here the parameter σ , one concludes on the functional dependence of three twistor components, i.e.

$$\Pi(G, \tau^1, \tau^2) = 0, \quad (19)$$

which, as in the real case (see (9)), demonstrates that fundamental null congruence of “rays” generated by the Newman’s charge is indeed *shear-free* for any its world line (or, differently, for any respective twistor function Π).

Let us recall now the fundamental property of *multivaluedness* of the GCRE solutions or, equivalently, – of the solutions to the SFC equations (9), or (17) in the considered case of the Newman’s congruence. Precisely, for any fixed point $Z = \{u, v, w, p\} \in \mathbb{CM}$ one obtains a countable (finite or infinite) set of solutions $\{G, \sigma\}$ of the system (17) so that, generically, *there exists a great number of continuous branches* – “modes” – of the twistor field functions $\mathbf{W}(Z)$, of the eikonal field $\sigma(Z)$, as well as a great number of respective shear-free *subcongruences*, superposing at each point Z . Notice that general concept of multivalued fields has been discussed in [2].

We see now that *any* complex ray of *any* mode of a SFC originates from a generating charge at some its position $\widehat{Z}(\sigma)$ which, therefore, is a *focal point* of the congruence. Thus, the World line of the Newman’s generating charge can be itself considered as the *focal line*.

One should distinguish between focal points with indefinite value of the main spinor field G and those of caustics (envelopes of the congruence rays) at which, generically, all of the field modes are well defined but two or more of them merge in one. For the Newman’s SFC caustic points, in account of (14), satisfy the condition

$$S' = \frac{dS}{d\sigma} = \widehat{u}'(v - \widehat{v}(\sigma)) + \widehat{v}'(u - \widehat{u}(\sigma)) - \widehat{w}'(p - \widehat{p}(\sigma)) - \widehat{p}'(w - \widehat{w}(\sigma)) = 0. \quad (20)$$

Locus of “hypercaustic” points – *cusps* – at which at least some *three* of the modes amalgamate – are analogously defined by the condition

$$S'' := \frac{d^2S}{d\sigma^2} = \widehat{u}''(v - \widehat{v}) + \widehat{v}''(u - \widehat{u}) - \widehat{w}''(p - \widehat{p}) - \widehat{p}''(w - \widehat{w}) - 2(\widehat{u}'\widehat{v}' - \widehat{w}'\widehat{p}') = 0 \quad (21)$$

and, finally, for the strongest singularities (*hypercusps*) one has the constraint $S''' = \dots = 0$. More stronger singularities have degenerate codimension space and generically exist in 4-dimensional $\mathbb{CM}$ space only as isolated points.

From (14) and (20) we easily find that at any focal point at least *two* modes merge together so that for any $Z = \widehat{Z}(\lambda)$, setting $\sigma = \lambda$, one gets identically $S = S' \equiv 0$. However, in the special case of a *null* World line defined as follows:

$$\det \|\widehat{Z}'\| \equiv \widehat{u}'\widehat{v}' - \widehat{w}'\widehat{p}' \equiv 0, \quad (22)$$

at any point $Z = \widehat{Z}(\lambda)$ on the World line with $\sigma = \lambda$ one can easily prove that more stronger conditions for singularity are satisfied: $S = S' = S'' = S''' = 0$, so that now *four* modes merge in one and are indefinite at the focal point itself. Further on we assume that *the World line of generating charge is null*.

Note that the considered equation of the null cone (14) in the case of the real space-time $\mathbf{M}$, in particular in the framework of electrodynamics, is known as the *retardation equation*. With real Cartesian coordinates $\{x, y, z, t\}$ and natural parametrization $\widehat{t}(s) = s = \Re(\sigma)$ it takes the form

$$c^2(t - s)^2 - (x - x(s))^2 - (y - y(s))^2 - (z - z(s))^2 = 0, \quad (23)$$

where the charge moves along a real world line $\{x(s), y(s), z(s)\}$. If the velocity v of the charge is assumed to be smaller than the light one, $v < c$, then equation (23) can be proved (see, e.g., [34]) to have always two roots only one from which has $s \leq t$ and satisfies thus the causality requirement. This fixes a unique position of the generating charge *in the past* which forms the field (electromagnetic Lienard-Wichert field in particular) at the observation point $\{x, y, z, t\}$ propagating to it rectilinearly with the speed of light. If the latter point belongs to the trajectory of the charge itself, equation (23) has only trivial solution $s = t$ (i.e., the retardation time is equal to zero, the observation point coincides with the “influence” one).

Situation changes drastically in the complex space-time. In fact, now a whole set of solutions $\{\sigma_N\}$ of the complex null cone (CNC) equation (14) does exist (for any point Z) which specify a (great) number of “influence points”, i.e. of the *images* of one and the same charge each forming its own mode of twistor field at Z and perceiving, therefore, by the observer at Z as a representative of the whole *ensemble of identical but distinguishable (via individual position and dynamics) point charges* – **duplicons**.

Notice that total *number* of duplicons does not depend, as a rule, on the observation point. However, a realistic “physical” observer $\mathbf{O}$ should be *material* and, therefore, in the simplest idealized case can be identified with a focal point itself. Then for the observation point one has $Z = \widehat{Z}(\lambda)$, with varying parameter $\lambda \in \mathbb{C}$ playing the role of the *complex proper time* of an “elementary” observer and manifesting itself as an evolutionary parameter (see below). Now four or two of the duplicons (the number depends on the fact is the World line null (22) or not) merge together forming the focal point of the observer $\mathbf{O}$ himself whereas the other $N - 4$ (or $N - 2$) duplicons are perceived by $\mathbf{O}$ as an ensemble of external identical point particles which are disposed and “move” with respect to $\mathbf{O}$ in the *observable 6D space-time* defined by the CNC equation (14). CNC geometry and its reduction to the 4D physical space-time will be studied in the next section.

3 Complex time and 6D physical geometry of the complex null cone

In the algebroid dynamics on the real $\mathbf{M}$ fundamental twistor field inevitably arises from the primary structure of $\mathbb{B}$ -differentiable functions-fields. By this, physical time t manifests itself as a *field evolutionary parameter*. Indeed, the incidence relation (7) is invariant under a 1-parameter group of translations of coordinates $X = X^+ \in \mathbf{M}$ along the rays of correspondent null congruence:

$$x_a \rightarrow x_a + n_a t, \quad (a = 1, 2, 3); \quad x_0 \rightarrow x_0 + t, \quad n_a = \xi^+ \sigma_a \xi / \xi^+ \xi, \quad \vec{n}^2 \equiv 1. \quad (24)$$

Presence of this symmetry demonstrates that the twistor field *propagates* from any initial point with universal velocity $v = c = 1$ along the directions specified by vector $\vec{n}$ (and locally defined by the field itself) or, equivalently, is *preserved* in value along these in the 4D-sense. Just the parameter t along the rays can be treated as local physical time (see correspondent discussion in [2, 24, 25]).

When one generalizes the scheme to the complex background, the same considerations lead to a more complicated and intriguing notion of time. Indeed, in $\mathbb{CM}$ corresponding symmetry group of translations is $2\mathbb{C}$ -parametric (the coordinates can vary across the so called *complex α -plane* [14]). We'll, however, start from an alternate representation of this field symmetry what for decompose the space $\mathbb{CM}$ of biquaternion algebra into two *real quaternionic* (4D Euclidean) spaces represented by 2×2 *unitary* matrices U and V :

$$Z = U + iV, \quad U^+ = U^{-1}, \quad V^+ = V^{-1}. \quad (25)$$

The first of these subspaces, say $\mathbf{U}$, may be considered as the main coordinate space whereas the second one $\mathbf{V}$ will then play the role of the *evolutional parameter space*. Precisely, from the incidence condition (13) and the decomposition (25) one gets, after some equivalent transformations, the following *translation law*:

$$U = U^{(0)} + V * N, \quad (26)$$

where the quaternion $U^{(0)}$ of initial coordinates is algebraically expressed via the twistor field components, quaternion $V = v_0 + iv_a \sigma_a$ defines four real evolutionary parameters $\{v_0, v_a\}$ ("quaternionic time"), and the *unit* quaternion $N = 1 + in_a \sigma_a$ (with the vector $\vec{n}$ of the same form as in the real case (24)) specifies local directions for the *translations preserving twistor field*.

Interestingly, the whole law (26) does not contain any residual of initial complex structure since all of the matrices therein are unitary and $(*)$ can be considered thus as multiplication in the *real quaternion algebra* $\mathbb{H}$. In the components, fundamental translation law (26) for four coordinates $\{x_\mu\}$, $\mu = 0, 1, 2, 3$ of the main *Euclidean* space $U \in \mathbf{E}^4$, $U = x_0 + ix_a \sigma_a$ has the following form:

$$\Delta x_a = n_a \Delta v_0 + \varepsilon_{abc} n_b \Delta v_c, \quad \Delta x_0 = n_a \Delta v_a, \quad (27)$$

so that any of translations satisfies one complex (two real) constraints $\Sigma(\Delta z_\mu)^2 = 0$, or in the components:

$$\begin{cases} (\Delta x_0)^2 + (\Delta x_1)^2 + (\Delta x_2)^2 + (\Delta x_3)^2 = (\Delta v_0)^2 + (\Delta v_1)^2 + (\Delta v_2)^2 + (\Delta v_3)^2 \equiv (\Delta t)^2, \\ (\Delta x_0)(\Delta v_0) + (\Delta x_1)(\Delta v_1) + (\Delta x_2)(\Delta v_2) + (\Delta x_3)(\Delta v_3) = 0, \end{cases} \quad (28)$$

fixing the structure of the *complex null cone*.

Notice that in the particular case when three of the parameters Δv_a vanish and the fourth one acquires the meaning of physical time $\Delta v_0 = t$, three “space” coordinates $\{x_a\}$ are translated just as in the former Minkowski case (24), i.e. rectilinearly and with fundamental “light” velocity whereas for additional Euclidean coordinate one has $\Delta x_0 = 0$, so that it is “frozen” and loses its dynamical sense.

Generically, the translation law (27) is much more complicated and rich in structure. The presence of four evolutionary parameters makes the field dynamics essentially uncertain since the *order* of continuous change of the parameters is not determined by the structure. However, we can naturally assume that the role of local physical time interval is now performed by the total Euclidean distance $(\Delta t)^2 = \Sigma(\Delta x_\mu)^2$ passing by the field in the coordinate space. In account of (28) this is equal to corresponding distance in the parameter space and twice smaller than that in the whole complex space with natural metric $\Sigma|z_\mu|^2$.

Thus, in the generic complex case local physical time can be related to the translations preserving the value of basic twistor field and to the whole distance to which the field propagates via these translations. In the latter aspect such treatment of physical time is similar to that proposed in the works of H.Montanus [26] and J.Almeida [27]. They considered the model of space-time as a 4D Euclidean space $\mathbf{E}^4$ with fourth coordinate being identified with local *proper time* of a particle of matter, and with local *coordinate time interval* being the metric interval covered by the latter (in natural units). By this, by analogy with the old ideas of F.Klein, Yu.Rumer et al., it is assumed that *all the material formations move in $\mathbf{E}^4$ with constant in modulus fundamental velocity $\Sigma(\Delta x_\mu/\Delta t)^2 = 1$ ($= c^2$)*. From here it follows immediately that *3-dimensional velocity of all material particles may be only less than or equal to the fundamental one*.

We see thus that in the context of successive biquaternionic electrodynamics, under the reduction of the primary “observable” geometry of the complex null cone of “elementary” observer, there arises a *model of the 4D Euclidean space-time* similar to that postulated in the approach of Montanus-Almeida. However, restriction of the 3-velocities’ values by the fundamental value holds in the algebrodynamics only for propagation of “prematerial” twistor field. As for the particle-like formations represented here by the focal and by the (hyper-) caustic points, this property will be identically fulfilled only for their *absolute* velocities, and only if the World line is null. For *relative* (“observable”) velocities situation is more complicated (see below). On the other hand, the structure of cone includes, apart of the main Euclidean “physical” space $\mathbf{U}$, a 2D orthogonal space $\mathbf{V}$ of the 2-sphere which could be treated (at least at the focal points of the particle’s location) as the space of directions of its *spin vector*.

Unfortunately, the scheme based on the concept of the 4D Euclidean space-time suffers from evident inconsistencies with Special Relativity. Indeed, though formally the first constraint in (28) can be represented in the form of the Minkowski-like interval $ds^2 := dx_0^2 =$

$dt^2 - dx_1^2 - dx_2^2 - dx_3^2$, there are no grounds for the proper time interval ds to remain invariant under the 4D rotations, as well as for the coordinate time dt to transform according to the Lorentz group. In the biquaternionic algebrodynamics situation is much more satisfactory since, under the transformations of the automorphism group $SO(3, \mathbb{C})$ of the $\mathbb{B}$ -algebra, the proper time represented by the zeroth coordinate, indeed remains invariant. Even by this, however, transformations of the spacial coordinates and coordinate time differ from the canonical Lorentz ones. This problem demands further investigation.

Let us recall now that in the algebrodynamical approach we have also to solve the above-mentioned problem of indefinite character of field evolution related to the presence of three additional free evolutionary parameters v_a or, on the other hand, – the problem of *ordering* of the 4D “quaternionic time”. This last difficulty can be partially overcome in a rather successive way. Note that under arbitrary field-preserving translations with four parameters $\{v_\mu\}$ the particle-like formations (focal points, caustics etc.) are not preserved and, generally, disappear from the main space $\mathbf{E}^4$.

In particular, if one takes all $v_\mu = 0$ to get the field-particle distribution at an *initial* instant of time, say at $t = 0$, then, generically, one will find the space $\mathbf{E}^4$ *empty*, i.e. free from any focal points. Indeed, the latter are *holomorphically* defined as the points of a complex World line $Z = \widehat{Z}(\sigma)$, and all four equations necessary for coordinates of these points to be *unitary* can’t be satisfied for any complex value of the parameter σ .

Thus, the problem is to concord the structure of the main quaternionic Euclidean space $\mathbf{U}$ and the “real quaternionic” type of the field evolution law (27) with the basic holomorphic structure of the null congruence and its particle-like singularities. To do this, let us return back to the above introduced *observable* space-time $\mathbf{O}$, i.e. to the 6D space-time of the complex null cone of an elementary observer $\mathbf{O}$ which is identified with a focal point and moves along its World line in synchro with the process of field evolution. On this *relative* subspace modelling physical space-time, we are now going to consider the evolutionary law for coordinates $\mathbf{U}$ preserving the structure of basic equations (14) and (20) together, i.e. *preserving both the value of twistor field and the condition of caustic*. This extended “field-caustic preserving” automorphism has the following $1\mathbb{C}$ -parameter holomorphic form:

$$Z \rightarrow Z + \tau(Z - \widehat{Z}(\sigma)), \quad \tau \in \mathbb{C}. \quad (29)$$

This means that caustics reproduce themselves along the complex straight lines ending (starting) at a focal point, and the parameter τ along these should be considered as the only physical evolutionary parameter – the “complex time”. In the real case such situation could correspond to a “signal” propagating rectilinearly with fundamental velocity. Complex structure, however, is not ordered, so there are infinitely many “ways” connecting some two points on such a line. Moreover, one is even not aware which of them is the emitter, and which – the receiver point. Note that the concept of two- (or multi- in general) dimensional time and the resulting problem of ordering of physical events have been considered in a number of works, in particular by Sakharov [28]. In the works of Yefremov [29, 30] just the structure of the automorphism group $SO(3, \mathbb{C})$ of the biquaternion algebra has been exploited, and at its base the concept of *three-dimensional time* was put forward. Under special orthogonality requirements this was assumed to reduce to the ordinary physical one-dimensional time.

Uncertainty of the field evolution process related to the complex nature of time is weaker than that for the primary “quaternionic time” but still too severe for usual physical interpretation, at least if one adheres to the paradigm of classical field theory.

To concord the concept of complex time with classical representations, one could assume, in addition to the holomorphic structure of the GCRE (or of the SFC equations), the existence of some “evolutionary curve” $\tau = \tau(s)$, $s \in \mathbb{R}$ on the complex plane which establishes the order of passing of different “states” in the main “physical” space $\mathbf{U}$. Such a curve as well as the World line of generating charge can be enormously complicated, possess the points of self-intersection etc. Because the observer has no information about the form of this curve, the future for him seems to be indefinite. However, a sort of *probabilities* can be introduced for different “continuations” of the evolutionary curve, and this procedure can open the way for explanation of quantum phenomena and for revealing the links of algebrodynamics and the Feynman’s version of quantum mechanics in particular. These problems will be discussed elsewhere.

There is evidently a local correspondence between the “evolutionary time” τ , the parameter σ of the World line and, say, the zeroth coordinate of the observer which can be treated as the “complex proper time”. In account of the quaternionic representation of the complex null cone of field evolution (28) we can interpret then the real part Δx_0 of “complex time increment” Δz_0 as an evolitional parameter in the main coordinate space $\mathbf{U}$ (“translational” proper time), whereas its *imaginary part* Δv_0 – as that in the orthogonal spin space (“rotational” proper time). Existence of the evolutionary curve implies also direct local connection between the increment of its parameter s and corresponding interval of the coordinate time Δt . Explicit formulas can be easily presented.

Let us consider now in more details the above mentioned process of emission/reception of a “signal-caustic” by “charges – focal points”. Let the “interacting” charges – duplicons $\hat{Z}(\lambda)$, $\hat{Z}(\sigma)$, $\sigma \neq \lambda$ are linked by the caustic. This corresponds to a solution of the joint system of equations (14) and (20), with substitution $Z = \hat{Z}(\lambda)$ instead of taking an arbitrary observation point Z . This system has a *discrete* set of pairs of solutions $\{\lambda, \sigma\}$ any of which correspond to two fixed *instants of complex time* of emission/reception of the caustic signal and, respectively, to two positions of the charges involved into the process.

By this, if the moment of observation λ “lies in the future” with respect to the “moment of influence” σ , i.e. corresponds to a greater value of the monotonically increasing parameter s of the evolutionary curve, than the observer $\hat{Z}(\lambda)$ would seem to *receive* the signal from the charge at $\hat{Z}(\sigma)$, and the latter would *emit* the signal towards $\hat{Z}(\lambda)$. The converse situation is evident. However, one should not forget that, in essence, this is *one and the same* charge *interacting with itself in its own past or future*.

Thus, all the *events* in complex space-time are well-defined and predetermined by the only structure of the World line of the generating charge (the focal line of the fundamental congruence). These are: the total number of charges – duplicons (“observed” by anyone of them; relative coordinates of duplicons *permanently* correlated with those of the observer through equal values of the primary twistor field); and, finally, a set of points on the World line at which this correlation between a pair of duplicons is strongly *amplified* (the charges are involved into the emission/reception process). However, the precise picture of evolution still remains indefinite and should be completed by additional information about the true form

of the evolutionary curve (extremely complicated and exceptional from the mathematical viewpoint).

A microscopic object (a focal point) is detectable for an idealized “elementary” observer only at some discrete moments of time at which the caustic line links them together. Near these, *relative* instantaneous velocity of the object is easily proved to be equal to its *absolute* one and, for the considered case of a *null* World line, is always less or equal to the fundamental velocity (the speed of light). These conclusions seem to hold also for a “classical” macroscopic object which is nearly permanently linked with a similar “classical” observer by a *net* of caustic lines. A detailed description (in the two utmost cases – quantum (microscopic) and classical (macroscopic)) of the correlated pairs “object - observer” exchanging by caustic “signals” will be presented elsewhere.

References

- [1] V.V.Kassandrov, in: *Physical Interpretation of Relativity Theory. Proc. Int. Sci. Meeting PIRT-2003*, eds. M.C.Duffy, A.N.Morozov, V.O.Gladyshev. – Moscow, Bauman Technical State Univ. Press, Liverpool, Sunderland, 2003, 23-34.
- [2] V.V.Kassandrov // *Hypercomplex Numbers in Geometry and Physics*, **1**, No.1, 2004, 91-107; (hep-th / 0312278).
- [3] V.V.Kassandrov, *The Algebraic Structure of Space-Time and the Algebrodynamics*. – Moscow, People’s Friendship Univ. Press, 1992 (in Russian).
- [4] V.V.Kassandrov // *Grav. & Cosm. (Moscow)*, **1**, No.3, 1995, 216-222; (gr-qc / 0007027).
- [5] V.V.Kassandrov // *Acta Applicandae Math.*, **50**, 1998, 197-206.
- [6] V.V.Kassandrov, in: *Has the Last Word been Said on Classical Electrodynamics?*, eds. A.Chubykalo, V.Onoichin, A.Espinoza, R.Smirnov-Rueda. – Rinton Press, 2004, 42-67; (physics / 0308045).
- [7] V.V.Kassandrov, V.N.Trishin // *Gen. Rel. Grav.*, **36**, 2004, 1603-1612; (gr-qc / 0401120).
- [8] V.V.Kassandrov, in: *Quasigroups and NonAssociative Algebras in Physics*, eds. Ja.K.Löhmus, P.Kuusk. – Tallinn, Inst. Phys. Estonia Proc., No.66, 202-212.
- [9] V.V.Kassandrov, J.A.Riscallah *E-print*, gr-qc / 0012109.
- [10] V.V.Kassandrov, J.A.Riscallah, in: *Geometrical and Topological Ideas in Modern Physics*, ed. V.A.Petrov. – Protvino, Inst. High Energy Phys., 2002, 199-212.
- [11] V.V.Kassandrov, J.A.Riscallah, in: *Recent Problems in Field Theory*, ed. A.V.Aminova. – Kasa, Kasa State Univ. Press, 1998, 176-186; (gr-qc / 9809078).

- [12] V.V.Kassandrov // *Vestnik Peopl. Friend. Univ., Physica*, **8(1)**, 2000, 34-45 (in Russian).
- [13] V.V.Kassandrov // *Grav. & Cosm. (Moscow)*, **8**, Suppl.2, 2002, 57-62; (math-ph / 0311006).
- [14] R.Penrose, W.Rindler *Spinors and Space-Time. Vol. II.* – Cambridge, Cambridge Univ. Press, 1986.
- [15] G.Debney, R.P.Kerr, A.Schild // *Journ. Math. Phys.*, **10**, 1969, 1842-1856.
- [16] V.V.Kassandrov, V.N.Trishin // *Grav. & Cosm. (Moscow)*, **5**, No.4, 1999, 272-276; (gr-qc / 0007026).
- [17] R.P.Kerr, W.B.Wilson // *Gen. Rel. Grav.*, **10**, 1979, 273-281.
- [18] B.Carter // *Phys. Rev.*, **174**, 1968, 1559-1572.
- [19] E.T.Newman // *Phys. Rev. D*, **65**, 2002, 104005-104018; (gr-qc / 0201055).
- [20] E.T.Newman // *Journ. Math. Phys.*, **14**, 1973, 102-105.
- [21] C.Kozameh, E.T.Newman, G.Silva-Ortigoza // *Class. Quant. Grav.*, **22**, 2005, 4679-4698; (gr-qc / 0506046).
- [22] E.T.Newman // *Class. Quant. Grav.*, **21**, 2004, 3197-3222; (gr-qc / 0402056).
- [23] R.W.Lind, E.T.Newman // *Journ. Math. Phys.*, **15**, 1974, 1103-1114.
- [24] V.V.Kassandrov, in: *Mathematics and Culture. Mathematics and Practice*, ed. M.Yu.Simakov. – M., “Self-Education”, 2001, p.61-76 (in Russian).
- [25] V.V.Kassandrov, in: *Proceedings of the Seminar on Study of Time Phenomenon*, ed. A.P.Levich. – M., “Binom”, 2006 (in print).
- [26] J.M.C.Montanus // *Hadron. Journ.*, **22**, 1999, 625-673.
- [27] J.B.Almeida // *E-print*, gr-qc / 0104029.
- [28] A.D.Sakharov // *Journ. Theor. Exper. Phys. (USSR)*, **87**, 1984, 375-383.
- [29] A.P.Yefremov // *Grav. & Cosm. (Moscow)*, **2**, № 1, 1996, 77-83.
- [30] A.P.Yefremov // *Grav. & Cosm. (Moscow)*, **2**, № 4, 1996, 335-341.
- [31] W.Kinnersley // *Phys. Rev.*, **186**, 1969, 1335-1336.
- [32] A.Ya.Burinskii // *Phys. Rev. D*, **67**, 2003, 12024-12040; (gr-qc / 0212048).
- [33] A.Ya.Burinskii // *Class. Quant. Grav.*, **20**, 2003, 905-912; (gr-qc / 0210010).
- [34] J.B.Jackson, *Classical Electrodynamics.* – N.Y., 1975, Ch. 14.